arXiv:quant-ph/0306153v1 23 Jun 2003

Why relativity allows quantum tunnelling to 'take no time'?

D.Sokolovski

School of Mathematics and Physics, Queen's University of Belfast, Belfast, BT7 1NN, United Kingdom

(February 1, 2008)

In quantum tunnelling, what appears an infinitely fast barrier traversal can be explained in terms of an Aharonov-like weak measurement of the tunnelling time, in which the role of the pointer is played by the particle's own coordinate. A relativistic wavepacket is shown to be reshaped through a series of subluminal shifts which together produce an anomalous 'superluminal' result.

PACS number(s): 03.65.Ta, 73.40.Gk

In 1932 MacColl's [1] noted that quantum tunnelling 'takes no time' in the sense that the tunnelled pulse can arrive in the detector ahead of its free counterpart, as if it has spent no time in the barrier region b. This effect been observed experimentally and extensively analysed by several authors (for a review see [2]). There is a broad consensus that what appears as 'superluminal' transmission takes place due to a reshaping mechanism (see, for example, [4]), but the role of the tunnelling time in the wavepacket propagation remain a contentious issue [3]. One dilemma one encounters when trying to explain the tunnelling speed up is whether quantum mechanics allows a particle to spend anomalously short ($\ll b/c$) durations in the barrier as was implied in [5] or, as we argue below, the effect can be achieved while maintaining, in agreement with special relativity, all velocities bounded by the speed of light c. A suggestion of how this may be possible, can be found in the work of Aharonov and coworkers [6–8], who have shown that under certain 'weak' conditions a von Neumann meter may produce readings well away from the region containing all eigenvalues of the measured variable. In this Letter, we demonstrate the analogy between wavepacket scattering and a weak measurement of the tunnelling time, which, although effectively all durations involved are subluminal, produces an apparently 'superluminal' result.

The authors of [6–8] considered a von Neumann measurement of a variable $\mathcal{A}$, $\mathcal{A}|\phi_i>=A_i|\phi_i>, 1\le i\le N,\quad N\gg 1,\quad 0\le A_i\le 1$ on a system which is prepared in a state $|I>=\sum_i a_i|\phi_i>$ and after the measurement is found (postselected) in another state $|F>=\sum_i b_i|\phi_i>$. After postselection, the pointer wavefunction in the coordinate representation is given by

$$<\tau|M>=\sum_{i=1}^{N} G(\tau-A_i)\eta_i, \qquad \eta_i\equiv b_i^* a_i \tag{1}$$

where $G(\tau)$ is the initial state of the meter. The 'paradox' observed by Aharonov *et al* is that if the measurement is weak (the width of $G(\tau)$ exceeds the separation between the eigenvalues A_i), a careful choice of $|I>$ and $|F>$ allows one to ensure that $<\tau|M>\approx G(\tau-\alpha)$ with an arbitrary shift α which can then be written as the weak value [6] of the operator $\mathcal{A}$,

$$\alpha=<F|\mathcal{A}|I>/<F|I>=\sum_{i=1}^{N}A_i\eta_i/\sum_{i=1}^{N}\eta_i. \tag{2}$$

Therefore it is possible (although improbable, since $|<F|I>|^2\ll 1$) that an experiment would produce, for example, a reading of 100 even though all eigenvalues of $\mathcal{A}$ involved do not exceed 1 [6]. In [8] this 'peculiar interference effect' is summarised as destructive interference in the whole "allowed" ($0<\tau<1$) region and the constructive interference of the tails in the "forbidden" ($\tau\approx 100$) region.

The relevance of the weak measurement analysis to wavepacket tunnelling, where no external meter is involved and no operator is available for the time spent in the barrier, is not obvious and will be established next. Classically, the amount of time a relativistic particle with a world line $x(s)=(x_0(s),\vec{x}(s))$ spends in a spatial region Ω is given by

$$t^{cl}_\Omega[x(s)]\equiv\int_{s'}^{s''}dsW_\mu(x)dx^\mu/ds=\int\theta_\Omega(\vec{x})dt. \tag{3}$$

where we have put, in the chosen frame of reference, $W_\mu(x)=(\theta_\Omega(\vec{x}),0,0,0)$ ($\theta_\Omega(\vec{x})\equiv 1$ for $\vec{x}\in\Omega$ and 0 otherwise). Quantally, Eq.(3) can be used to distinguish between scenarios in which a particle spends different durations inside Ω. Propagator for a spin-zero Klein-Gordon particle in an electromagnetic field $A_\mu(x)$ can be written as a sum over four-dimensional paths parametrised by the fifth parameter u, [9,10] ($\partial_z\equiv\partial/\partial z, \hbar=c=1$)

$$g(x,x')=\int_0^\infty du\exp(-im^2u/2)\int Dx\exp[-i\int_0^u\mathcal{L}(x)du] \tag{4}$$

where $\mathcal{L}(x)=(\partial_u x^\mu)^2/2+e\partial_u x^\mu A_\mu$. With the help of Eqs.(3) and (4) the propagator restricted only to those paths which spent inside Ω a duration τ, $\int Dx\to\int Dx\delta(t^{cl}_\Omega[x(u)]-\tau)$ is seen to be given by

$$g(x,x'|\tau)=(2\pi)^{-1/2}\int_{-\infty}^{\infty}\exp(iW\tau)g(x,x'|W)dW \tag{5}$$

where $g(x,x'|W)$ is the propagator for the modified electromagnetic field $A_0(x)+W\theta_\Omega(\vec{x}),\vec{A}(x)$. Analysing the time evolution of $g(x,x'|\tau)$ leads to the clocked Klein-Gordon equation ($p_\mu \equiv -i\partial_{x_\mu}$)

$$\{[i\partial_t + i\partial_\tau\theta_\Omega(\vec{x}) - eA_0]^2 - (\vec{p} - e\vec{A})^2 - m^2\}\Phi(x|\tau) = 0 \quad (6)$$

where $\Phi(x|\tau)$ is the (scalar) amplitude that a particle at a space-time point x has previously spent a duration τ in the region Ω. In the case of the Dirac electrons, no path integral is available for the spin degrees of freedom. However, one can still define a (4x4 matrix) amplitude F_P for a given space-time path [9]. On application of a field $A(x)$ the phase of F_P is modified, $F_P^A = \exp\{-i\int A_\mu dx^\mu\}F_P$, and restricting the paths as above yields the clocked Dirac equation

$$\{[i\partial_t + i\partial_\tau\theta_\Omega(\vec{x}) - eA_0]\gamma_0 - (\vec{p} - e\vec{A})\vec{\gamma} - m\}\Phi(x|\tau) = 0. \quad (7)$$

Note that like the clocked Schrödinger equation [21], both Eqs.(6) and (7) are obtained from their original versions by replacing $i\partial_t \to i\partial_t + i\partial_\tau\theta_\Omega(\vec{x})$.



FIG. 1. Transmission of a Gaussian wavepacket across the region $[-b/2, b/2]$. Inset: $|\Psi^T(z,t)|$ for free motion and tunnelling ($\times A \equiv 10^{-2}T(p_0,V)$) for $\epsilon_0\tau_c/\hbar = 6000$; $\epsilon_{p_0}\tau_c/\hbar = 6007.5$; $V\tau_c/\hbar = 15.$; $\Delta z/b = 0.55$; $z_0/b = 2.5; t/\tau_c = 100.$; $\tau_c \equiv b/c$ Also shown by the dashed line is the semiclassical result Eq.(18). (a) the factor $G(z + z_0 - b - v_0(t-\tau))$ in Eq.(13); Real part of $\eta(p_0,t)$ for (b) free motion, $V = 0$; and (c) tunnelling, $V\tau_c/\hbar = 15$.

To study the durations involved in in one-dimensional transmission of a spinless particle across a rectangular potential barrier $V\theta_{-b/2,b2}(z)$ considered in [12], we prepare a wavepacket with a mean momentum p_0 and a momentum (coordinate) space width Δp (Δz) initially containing only negative frequencies (no antiparticles)

$$\Psi(z,t=0) = \int A(p)exp(ipz)dp = exp(ip_0z)G(z+z_0). \quad (8)$$

where $z_0 \gg b/2$ and a suitable $G(z)$ (e.g., a Gaussian $G(z) = exp(-z^2/\Delta z)$) is chosen to ensure that all momenta in (8) tunnel. Solving Eq.(6) with the initial condition $\Phi(z,t=0|\tau) = \Psi(z,t=0)\delta(\tau)$ yields, for sufficiently large z and t, the traversal time decomposition of the transmitted part of the wavepacket,

$$\Phi(z \gg b/2, t|\tau) = \int dpA(p)exp(ipz - i\epsilon_p t)\eta(p,\tau) \quad (9)$$

$$\eta(p,\tau) \equiv (2\pi)^{-1/2}\exp(-iV\tau)\int_{-\infty}^{\infty}\exp(iW\tau)T(p,W)dW \quad (10)$$

where $\epsilon_p = (p^2+m^2)^{1/2}$ and $T(p,W)$ is the transmission amplitude for the rectangular potential $W\theta_{-b/2,b2}(z)$. The amplitude $\Phi(z,t|\tau)$ gives the transmitted field under the condition that a particle spend in the barrier exactly a duration τ and a coherent sum over all such durations produces the transmitted pulse $\Psi^T(z,t)$

$$\Psi^T(z,t) = \int_{-\infty}^{\infty}\Phi(z,t|\tau)d\tau = \int T(p,V)A(p)exp(ipz - i\epsilon_p t)dp \quad (11)$$

For tunnelling, ($\epsilon_p - m < V \ll m$), $\Psi^T(x,t|V)$ although greatly reduced, lies approximately a distance b ahead of the free ($V = 0$) pulse suggesting that the particle has traversed the barrier instantaneously (see inset in Fig.1). To see how this speed up is achieved, we consider the semiclassical limit $pb, kb \gg 1$ [14], replace $T(p,W)$ by the first term in its multiple scattering expansion [13] ($k(W) \equiv [(\epsilon_p - W)^2 - m^2]^{1/2}$)

$$T(p,W) \approx \frac{4pk}{(p+k)^2}exp[ikb - ipb], \quad (12)$$

which is permissible both well above, $V \ll \epsilon_p - m$, and well below, $V \gg \epsilon_p - m$, the barrier, and evaluate the resulting integrals asymptotically. This yields

$$\Psi^T(z,t) \approx \exp(ip_0z - i\epsilon_{p_0}t) \times \int_{-\infty}^{\infty}G(z + z_0 - b - v_0(t-\tau))\eta(p_0,\tau)d\tau \quad (13)$$

together with the saddle point result for $\eta(p_0,\tau)d\tau$ [15] ($\hbar$ and c are reintroduced)

$$\eta(p_0,\tau) = \frac{(2\pi\hbar)^{-1/2}4p_0b\epsilon_0^{3/2}\tau_c^3}{(\tau^2-\tau_c^2)^{1/4}[p_0b(\tau^2-\tau_c^2)^{1/2} + \epsilon_0\tau_c^2]^2} \times exp\{\frac{i}{\hbar}[(\epsilon_p - V)\tau - \epsilon_0(\tau^2-\tau_c^2)^{1/2}] - i\pi/4\} \quad (14)$$

where $v_0 \equiv \partial\epsilon_p/\partial p_0$ and we have introduced $\epsilon_0 \equiv mc^2$ and $\tau_c \equiv b/c$. Since most of our derivation relies on the analytical properties of the wavevector $k(W)$, a similar analysis can also be applied to the Dirac electrons.

We proceed with the details of the reshaping mechanism responsible for the speed up of the tunnelled pulse. The first term in the integrand of Eq.(13) is the wavepacket translated as a whole along a classical trajectory which spends inside $[-b/2, b/2]$ a time τ and has the velocity

v_0 outside the region (Fig.1a) so that the total distance covered by the time t is

$$z - z_0 = b + v_0(t - \tau). \tag{15}$$

Each wavepacket is weighted by the (complex valued) amplitude $\eta(p_0, \tau)$ for a transmitted particle whose momentum is exactly p_0 to spend the duration τ in $[-b/2, b/2]$. It is instructive to consider the case of free motion first. For $V = 0$, the oscillatory amplitude $\eta(p_0, \tau)$ shown in Fig.1b coincides for $\tau \gg \tau_c$ with the non-relativistic result of [13] and has a stationary region around the classical value b/v_0. For $|\tau| < \tau_c$, it rapidly decays so that, owing to the relativistic nature of Eq.(6), the durations which imply superluminal velocities $v_{bar} = b/\tau$ are explicitly absent from the integral (13). For $\tau < -\tau_c$, $\eta(p_0, \tau)$ becomes oscillatory again due to the relativistic paths in Eq.(4) which experience time reversals associated with creation and annihilation of particle-antiparticle pairs [9,10]. The frequency of these oscillations is extremely high and, as one would expect, the processes involving virtual pair production do not contribute to the propagation. As a result, $\eta(p_0, \tau)$ selects the shape $G(z - z_0 - v_0 t)$ corresponding to the classical value $\tau = b/v_0$ and, in accordance with classical mechanics, one finds the initial pulse translated without distortion along the classical path $z(t) = v_0 t + z_0$.

In the case of tunnelling, the amplitude $\eta(p_0, \tau)$ differs from its free counterpart by the factor $\exp(-iV\tau/\hbar)$ in Eq.(10) which destroys the stationary region on the real τ-axis (Fig 1c). There is no preferred duration describing tunnelling and the analogy with Aharonov's weak measurement is now evident. Just as in Eq.(1) a sum of Gaussians all shifted by no more than 1 produces a Gaussian with an arbitrary large shift α, a continuos superposition of the copies of the initial wavepacket all propagated with purely subluminal velocities produces a pulse which looks as if it had spent a zero time in the barrier. The role of the pointer position is played by the particle's coordinate z, and the two distinctive features of a weak measurement are in place: the particle is postselected in a highly unlikely (most particles are reflected) transmitted state and the accuracy, which is determined by the coordinate width of the wavepacket, Δz must remain poor in order to prevent the particle passing over the barrier, which would destroy the effect. Several known properties of the tunnelling transmission can now be clarified. Destructive interference between the maxima and the backward tails of all Gaussians in Fig.1a explains why, as observed in [12], the transmitted pulse is formed from the front part of the initial wavepacket. Also, like a weak von Neumann meter [7], quantum tunnelling cannot be used to transmit information which cannot be extracted without a reference to 'superluminal' behaviour. Indeed, truncating G(z) after some $z' >> \Delta z$ would guarantee effective vanishing of $\Psi^T(z, t)$ in Eq.(13) for $z > ct - z_0$. Thus, rather than facilitating faster-than-light information transfer, a barrrier acts as a filter [7], rearranging the amplitudes in the forward tails of all (subluminally shifted) G's in Eq.(11) into a single peak.

Previous work [5,17–19] on relating the tunnelling time problem to weak measurements focused mostly on interpreting the 'complex time' [20–22]

$$\bar{\tau} = \int \tau \eta(p, \tau) d\tau / \int \eta(p, \tau) d\tau \tag{16}$$

as the weak value (2) of the projector onto the barrier region, $\theta_{[-b/2,b/2]}(z)$ [17,18] or its time average, $T^{-1} \int_0^T \theta_{[-b/2,b/2]}(t) dt$ [19]. In [5] the authors applied a low accuracy Salecker-Wigner clock to Dirac electrons obtaining, effectively, an estimate $v'_{bar} = b/Re\bar{\tau}$ for the mean velocity in the barrier. For a relativistic particle v'_{bar} may exceed c even though $\eta(p_0, \tau)$ has no support inside $-b/c < \tau < b/c$ and [5] questioned the significance of this result for predicting 'superluminal' transmission of electrons. The answer is that $\bar{\tau}$, which describes the work of an inaccurate external clock [17,21,22], cannot, in general, predict the position of the tunnelled pulse, whose shape is determined by the oscillatory integral (11) involving all moments of $\eta(p_0, \tau)$ for a range of particle's momenta p. The semiclassical case (11)-(14) is special because $\eta(p_0, \tau)$ has an imaginary saddle at

$$\tau_V = -i\tau_c \frac{(\epsilon_{p_0} - V)}{[\epsilon_0^2 - (\epsilon_{p_0} - V)^2]^{1/2}}, \tag{17}$$

and the superposition of purely subluminal shifts (11) results for G in an imaginary shift by $\alpha = -i|\tau_V|$. thus, within our approximation (see inset in Fig.1),

$$\begin{aligned} \Psi^T(z, t) \approx\, & G(z + z_0 - b - v_0(t + i|\tau_V|)) \\ & \times T(p_0, V) \exp[i(p_0 z - \epsilon_{p_0} t)/\hbar] \end{aligned} \tag{18}$$

and $\bar{\tau} \approx -i|\tau_V| + O(\hbar)$ is the single parameter governing the shape of the pulse and, therefore, the speed of transmission. As the tunnelling speed up effect itself, anomalously small values of $Re\bar{\tau}$ have their origin in the oscillatory nature of the traversal time amplitude distribution $\eta(p_0, \tau)$, but, in the full quantum case, the condition $\bar{\tau} < b/c$ is not sufficient for the 'superluminal' transmission to take place and a more extensive analysis based on Eq.(11) must is necessary.

In summary, special relativity restricts the range of tunnelling times to the subluminal domain, yet interference accounts for the 'superluminal' appearance of the transmitted pulse. This somewhat counterintuitive effect is most easily explained in terms of Aharonov's weak measurements. A barrier reshapes a semiclassical wavepacket by propagating components of the incident pulse with purely subluminal velocities before reassambling them with complex weights determined by the potential. Examining the position of the tunnelled particle, one may estimate the time it has spent in the barrier. Of necessity, the accuracy of such estimate (determined by the initial spread in the particle's position) is so poor that it yields an anomalously small 'forbidden' value, suggesting that the barrier was traversed almost instantaneously, just as a weak measurement on a spin 1/2 may return the value 100 [6]. This analysis differs from that of Refs. [6–8] in that it, contrary to the claim that the traversal time cannot be defined 'intrinsically' [23], involves no external pointer. That the role of such pointer can be played by the particle's own coordinate, can be traced back to the classical relation (15) between the distance and the time spent in the region, which below the barrier takes the form of weak measurement-like correlations between the two quantities. Finally we note that, given the universal nature of the weak measurement analysis stressed in [6–8,17], it is unlikely that a deeper or more 'classical' explanation could be found for the apparent superluminality observed in quantum tunnelling.

---

[1] L.A. MacColl, Phys.Rev. **40**, 621 (1932)

[2] R.Y. Chiao and A.M. Steinberg, Prog. Opt. **37**, 345 (1997)

[3]) *Time in Quantum Mechanics*, edited by J.G. Muga, R. Sala-Mayato

[4] F.E.Low, Ann.Phys.(Leipzig),**7**, 660 (1998) and I.L. Eguisquiza (Springer, 2002)

[5] C.R.Leavens and R.Sala Mayato, Ann.Phys.(Leipzig),**7**, 662 (1998)

[6] Y. Aharonov, D.Z. Albert and L. Vaidman, Phys. Rev. Lett. **60**, 1351 (1988)

[7] Y. Aharonov and L. Vaidman, Phys.Rev.A **41**, 11 (1990)

[8] Y. Aharonov and L. Vaidman, in [3], pp. 369-412

[9] R.P. Feynman, Phys.Rev. **80**,440 (1950)

[10] L.S. Schulman, *Techniques and Applications of Path Integration* (Wiley, New York, 1981), pp.225-236

[11] D. Sokolovski, Phys.Rev.Lett., **79**,4946 (1997)

[12] J.M. Deutch and F.E.Low, Ann.Phys.**228**, 184 (1993)

[13] D.Sokolovski and Y.Liu, Phys.Rev.A **63**, 012109 (2000)

[14] We also choose the momentum space width of the wavepacket Δp to be sufficiently small, $b\Delta p$, $b\partial k/\partial p_0 \Delta p$, $p_0 \Delta p t \approx 1$ and neglect terms $\approx (\Delta p)^2$.

[15] The branch of $(\tau^2 - \tau_c^2)^{1/2}$ is chosen to be positve below the right cut in the complex τ-plane with cuts along $[-\infty, \tau_c]$ and $[\tau_c, \infty]$. The contour of intregration in Eq.(11) runs above the left and below the right cut.

[16] D. Sokolovski, in [3], pp. 183-216

[17] A.M. Steinberg, Phys.Rev.Lett., **74**,2405 (1995)

[18] A.M. Steinberg, Phys.Rev.A, **52**,32 (1995)

[19] G. Iannaccone, in *Time and Its Implications*, edited by D. Mugnai, A. Ranfagni and L.S. Schulman (World Scientific, 1997), pp. 292-309

[20] H. Fertig, Phys.Rev.Lett., **65**,2321 (1995)

[21] D. Sokolovski, Phys.Rev.Lett., **79**,4946 (1997)

[22] D. Sokolovski, in [3], pp. 183-216

[23] N. Yamada, Phys.Rev.Lett.,**17**, 3350 (1999); a detailed response to Yamada's critique can be found in D.Sokolovski, Phys.Rev.A, 2002 (in press).